VOL. 512 JOURNAL OF CHROMATOGRAPHY JULY 20, 1990

CONTENTS

9TH INTERNATIONAL SYMPOSIUM ON HPLC OF PROTEINS, PEPTIDES AND POLYNUCLEOTIDES, PHILADELPHIA, PA, NOVEMBER 6–8, 1989

Foreword
by M. T. W. Hearn (Clayton, Australia) 1

Binding of sodium dodecyl sulphate to an integral membrane protein and to a water-soluble enzyme. Determination by molecular-sieve chromatography with flow scintillation detection
by M. Wallstén and P. Lundahl (Uppsala, Sweden) 3

Systematic optimization of protein separations on high-performance ion-exchange chromatographic media
by W. Warren, N. E. Astephen and T. E. Wheat (Milford, MA, U.S.A.) 13

Evaluation of factors which affect column performance with immobilized monoclonal antibodies. Model studies with a lysozyme–antilysozyme system
by M. T. W. Hearn and J. R. Davies (Clayton, Australia) 23

High-performance liquid chromatography of amino acids, peptides and proteins. XCVIII. The influence of different displacer salts on the bandwidth properties of proteins separated by gradient elution anion-exchange chromatography
by A. N. Hodder, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 41

High-performance liquid chromatography of amino acids, peptides and proteins. XCIX. Comparative study of the equilibrium refolding of bovine, porcine and human growth hormone by size-exclusion chromatography
by M. Fridman, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) 57

Predicting the performance of gel-filtration chromatography of proteins
by S. Yamamoto, M. Nomura and Y. Sano (Ube, Japan) 77

Preparative chromatography of proteins. Design calculation procedure for gradient and stepwise elution
by S. Yamamoto, M. Nomura and Y. Sano (Ube, Japan) 89

High-performance liquid chromatography of amino acids, peptides and proteins. CIII. Mass transfer resistances in ion-exchange and dye-affinity chromatography of proteins
by A. Johnston and M. T. W. Hearn (Clayton, Australia) 101

Ion-exchange high-performance liquid chromatographic isolation of the major allergen of parietaria pollen extract
by I. Søndergaard (Copenhagen, Denmark) and P. Falagiani, E. Bolzacchini, P. Civaroli, G. Di Gregorio, V. Madonini, A. Morelli and B. Rindone (Milan, Italy) 115

High-performance liquid chromatographic approach to the separation of antiviral and immunostimulant fractions in Neuramide
by N. Miraglia and B. Rindone (Milan, Italy), G. Folchitto (Ariccia, Italy), P. Amicucci and G. Antonelli (Rome, Italy) and A. Lanfranchi and M. Massa (Pavia, Italy) 123

Purification of a amylase–pullulanase bifunctional enzyme by high-performance size-exclusion and hydrophobic-interaction chromatography
by C.-H. Kim and D.-S. Kim (Tokyo, Japan), H. Taniguchi (Tsukuba, Japan) and Y. Maruyama (Tokyo, Japan) 131

Purification of immunomodulatory factors in human peripheral blood leukocytes
by P. C. Montecucchi, R. Caldarini, M. Sansó and A. M. Bertolaso (Cassina de' Pecchi, Italy) and V. E. M. Rosso Di San Secondo, G. Piccolo and G. Sirchia (Milan, Italy) 139

Oligosaccharide mapping of therapeutic glycoproteins by high-pH anion-exchange high-performance liquid chromatography
by R. Kumarasamy (Kenilworth, NJ, U.S.A.) 149

Isolation of isoproteins from monoclonal antibodies and recombinant proteins by chromatofocusing
by A. Jungbauer, C. Tauer, E. Wenisch, K. Uhl, J. Brunner, M. Purtscher, F. Steindl and A. Buchacher (Vienna, Austria) 157

Separation of fragments from human serum albumin and its charged variants by reversed-phase and cation-exchange high-performance liquid chromatography
by P. Iadarola, M. C. Zapponi, L. Minchiotti, M. L. Meloni, M. Galliano and G. Ferri (Pavia, Italy) 165

Immobilized metal ion affinity chromatography for the purification of *Fel d* I, a cat major allergen, from a house-dust extract
by J.-P. Dandeu, J. Rabillon, M.-J. Beltrand, M. Lux, R. Duval and B. David (Paris, France) 177

Chromatography and generation of specific antisera to synthetic peptides from a protective *Boophilus microplus* antigen
by P. J. Sharp, B. V. McInerney and D. R. J. Smith (Roseville, Australia), I. F. Turnbull (Parkville, Australia), D. H. Kemp (Indooroopilly, Australia) and K. N. Rand and G. S. Cobon (Roseville, Australia) 189

Characterization of apolipoproteins from chicken plasma
by E. M. Brown and H. J. Dower (Philadelphia, PA, U.S.A.) 203

Rapid high-performance liquid chromatographic protein quantitation of purified recombinant Factor VIII containing interfering substances
by W. Schröder (Wuppertal, F.R.G.) and M. L. Dumas and U. Klein (Berkeley, CA, U.S.A.) 213

Purification of membrane protein complexes isolated from a cyanobacterial thylakoid membrane by high-performance liquid chromatography
by M. Rögner (Wilmington, DE, U.S.A.) 219

Simultaneous determination of *Bordetella pertussis* toxin and filamentous haemagglutinin concentrations by hydroxyapatite high-performance liquid chromatography
by P. Chong, G. Jackson, W. Cwyk and M. Klein (Willowdale, Canada) 227

High-performance metal chelate interaction chromatography of proteins with silica-bound ethylenediamine-N,N′-diacetic acid
by M. D. Bacolod and Z. El Rassi (Stillwater, OK, U.S.A.) 237

High-performance liquid chromatographic investigations on the cleavage kinetics of side-chain-protected arginine derivatives with a sophisticated post-column reaction detector
by G. Grübler, F. Gutjahr, H. F. Echner, W. Voelter and H. Bauer (Tübingen, F.R.G.) . 249

Identification of peptides containing aromatic amino acids, cysteine, iodotyrosine and iodothyronine by high-performance liquid chromatography with photodiode-array detection
by J. Escribano, M. Asunción, J. Miguel, L. Lamas and E. Méndez (Madrid, Spain) . . . 255

Reversed-phase high-performance liquid chromatographic separation of synthetic phosphopeptide isomers
by L. Otvos, Jr., I. A. Tangoren and K. Wroblewski (Philadelphia, PA, U.S.A.), M. Hollosi (Budapest, Hungary) and V. M.-Y. Lee (Philadelphia, PA, U.S.A.) 265

Optimization of high-performance liquid chromatographic peptide separations with alternative mobile and stationary phases
by P. M. Young and T. E. Wheat (Milford, MA, U.S.A.) 273

Analysis of amino acids by liquid chromatography after pre-column derivatization with 4-nitrophenylisothiocyanate
by S. A. Cohen (Milford, MA, U.S.A.) 283

High-performance liquid chromatographic separation and partial characterization of the α- and β-chains of Alaskan sockeye salmon hemoglobin
by L. K. Duffy and R. A. Reynolds (Fairbanks, AK, U.S.A.) and J. P. Harrington (Mobile, AL, U.S.A.) 291

High-performance displacement chromatography–mass spectrometry of tryptic peptides of recombinant human growth hormone
by J. Frenz, J. Bourell and W. S. Hancock (San Francisco, CA, U.S.A.) 299

Tryptic fingerprinting on a poly(styrene–divinylbenzene) reversed-phase column
by J. K. Swadesh (Amherst, MA, U.S.A.) 315

Peptide mapping of HIV polypeptides expressed in *Escherichia coli*. Quality control of different batches and identification of tryptic fragments containing residues of aromatic amino acids or cysteine
by S. Renlund and I.-M. Klintrot (Uppsala, Sweden), M. Nunn and J. L. Schrimsher (La Jolla, CA, U.S.A.) and C. Wernstedt and U. Hellman (Uppsala, Sweden) 325

Synthetic antibody fragment as ligand in immunoaffinity chromatography
by G. W. Welling, T. Geurts, J. van Gorkum, R. A. Damhof, J. W. Drijfhout (Groningen, The Netherlands), W. Bloemhoff (Leiden, The Netherlands) and S. Welling-Wester (Groningen, The Netherlands) 337

Immobilization of Protein A at high density of azlactone-functional polymeric beads and their use in affinity chromatography
by P. L. Coleman, M. M. Walker, D. S. Milbrath, D. M. Stauffer, J. K. Rasmussen, L. R. Krepski and S. M. Heilmann (St. Paul, MN, U.S.A.) 345

Preparative high-performance liquid chromatography on a unique high-speed macroporous resin
by L. L. Lloyd and F. P. Warner (Church Stretton, U.K.) 365

Binding of lysozyme on the surface of entrapped phosphatidylserine–phosphatidylcholine vesicles and an example of high-performance lipid vesicle surface chromatography
by Q. Yang and P. Lundahl (Uppsala, Sweden) 377

Isolation of anti-idiotypic antibodies by immunoaffinity chromatography on Affinichrom beads
by T. M. Phillips (Washington, DC, U.S.A.) and J. V. Babashak (Vineland, NJ, U.S.A.) . 387

Assay of cytosine and cytidine deaminases by means of reversed-phase high-performance liquid chromatography
by P. K. Dutta, M. S. Shanley and G. A. O'Donovan (Denton, TX, U.S.A.) 395

Characterization of temperature-sensitive cytidine triphosphate synthase mutations in bacteria by high-performance liquid chromatography
by A. J. Bailey, P. K. Dutta, M. S. Shanley and G. A. O'Donovan (Denton, TX, U.S.A.) 403

Preparation of ethylenediaminephosphoramidates of nucleotides and derivatization with fluorescein isothiocyanate
by A. N. Al-Deen (Alexandria, Egypt), D. C. Cecchini and S. Abdel-Baky (Boston, MA, U.S.A.), N. M. A. Moneam (Alexandria, Egypt) and R. W. Giese (Boston, MA, U.S.A.) . 409

Convenient purification of tritylated and detritylated oligonucleotides up to 100-mer
by T. L. Hill and J. W. Mayhew (San Ramon, CA, U.S.A.) 415

Rapid separation, quantitation and purification of products of polymerase chain reaction by liquid chromatography
by E. D. Katz and L. A. Haff (Norwalk, CT, U.S.A.) and R. Eksteen (Woburn, MA, U.S.A.) 433

Author Index 445

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)